

## INFORM UC: An Update on Contemporary Management of UC



### **Maria T. Abreu, MD**

Professor of Medicine  
Professor of Microbiology and Immunology  
Martin Kalser Chair in Gastroenterology  
Director, Crohn's & Colitis Center  
University of Miami  
Miller School of Medicine  
Miami, Florida



### **Stephen B. Hanauer, MD, FACG**

Clifford Joseph Barborka Professor  
of Medicine  
Medical Director, Digestive Health Center  
Northwestern University Feinberg School  
of Medicine  
Chicago, Illinois



### **Gary R. Lichtenstein, MD, FACP, AGAF, FACP**

Director, Center for Inflammatory Bowel Disease  
The University of Pennsylvania Health System  
Professor of Medicine  
The Raymond and Ruth Perelman School of  
Medicine of the University of Pennsylvania  
Philadelphia, Pennsylvania

Accredited by Rehoboth McKinley  
Christian Health Care Services



Provided by the Gi Health Foundation



Supported by an educational grant from  
Pfizer, Inc.

**A CME Activity**  
Approved for  
1.0 AMA PRA  
Category 1 Credit™

**Release Date:** August 2020

**Expiration Date:** August 31, 2021

**Estimated time to complete activity:** 1.0 hour

**ON THE WEB:**  
[gastroenterologyandhepatology.net](http://gastroenterologyandhepatology.net)

Indexed through the National Library of Medicine  
(PubMed/Medline), PubMed Central (PMC), and EMBASE

## INFORM UC: An Update on Contemporary Management of UC

To claim 1.0 *AMA PRA Category 1 Credit*<sup>™</sup>  
for this activity, please visit:  
[gihealthfoundation.org/EVAL-INFORMUC](http://gihealthfoundation.org/EVAL-INFORMUC)

### Goal Statement

The primary goal of this CME monograph is to encourage the application of the latest advances in evidence-based medicine to improve the quality of care and improve outcomes for patients affected by UC.

### Target Audience

This CME monograph will target gastroenterologists, primary care physicians, nurse practitioners, physician assistants, and nurses.

### Educational Objectives

Upon completion of this educational activity, participants should be able to:

- Discuss key data regarding the latest diagnostic and treatment strategies for UC presented at DDW 2020
- Incorporate the latest advances in UC management into clinical practice
- Summarize implications of the current coronavirus pandemic on the management of patients with UC

### Accreditation Statement and Credit Designation

This activity has been planned and implemented in accordance with the accreditation requirements and policies of the New Mexico Medical Society (NMMS) through the joint providership of Rehoboth McKinley Christian Health Care Services (RMCHCS) and the Gi Health Foundation. RMCHCS is accredited by the NMMS to provide continuing medical education for physicians.

RMCHCS designates this activity for a maximum of 1.0 *AMA PRA Category 1 Credit*<sup>™</sup>. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Supported by an educational grant from Pfizer, Inc.

Provided by the 

Accredited by 

### Disclosures

Faculty members are required to inform the audience when they are discussing off-label, unapproved uses of devices and drugs. Physicians should consult full prescribing information before using any product mentioned during this educational activity.

#### Maria T. Abreu, MD

Speaker: Janssen, Prometheus Biosciences, Takeda, Focus Medical Communications, Boehringer Ingelheim Pharmaceuticals, Gilead, Imedex, Cornerstone Health, Inc, Landos Biopharma, UCB Biopharma SRL, Eli Lilly, and Cosmo Biopharma

#### Stephen B. Hanauer, MD, FACP

Speaker: AbbVie, Janssen, Pfizer, and Takeda; Consultant: AbbVie, Allergan, Amgen, Arena, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, Celltrion, Genentech, Gilead, GSK, Janssen, Lilly, Merck, Nestle, Novartis, Pfizer, Progenity, Prometheus, Receptos, Salix, Samsung Bioepis, Seres Therapeutics, Takeda, TiGenex, UCB Pharma, and VHSquared

#### Gary R. Lichtenstein, MD, FACP, AGAF, FACP

AbbVie (Consultant), American College of Gastroenterology (Honorarium for Associate Editor of *American Journal of Gastroenterology*), Celgene (Research, Consultant), CellCeutix (Consultant), *Clinical Advances in Gastroenterology* (Associate Editor–Honorarium), Ferring (Consultant), *Gastroenterology & Hepatology* [Gastro-Hep Communications] (Editor–Honorarium), Gilead (Consultant), Janssen Ortho Biotech (Consultant, Research, Funding to University of PA [IBD Fellow Education]), Eli Lilly (Consultant, DSMB), Luitpold/American Regent (Consulting, Honorarium [CME Programs]), Merck (Consulting, Honorarium [CME Program]), McMahon Publishing (Author [Honorarium]), Pfizer Pharmaceuticals (Consultant, Research, Funding to University of PA [IBD Fellow Education]), Prometheus Laboratories, Inc (Consultant), Romark (Consultant, Honorarium for CME), Salix Pharmaceuticals/Valeant (Consultant, Research), Shire Pharmaceuticals (Consultant, Research), SLACK, Inc (Book Royalty), Springer Science+Business Media (Editor–Honorarium), Takeda (Consultant, Funding to University of PA [IBD Fellow Education]), UCB (Consultant, Research), and Up-To-Date (Author–Honorarium)

**Julianne Messick, PharmD, Medical Writer** No real or apparent conflict of interest

Indexed through the National Library of Medicine (PubMed/Medline), PubMed Central (PMC), and EMBASE

#### Disclaimer

This supplement is supported by an educational grant from Pfizer, Inc. Support of this supplement does not imply the supporter's agreement with the views expressed herein. Every effort has been made to ensure that drug usage and other information are presented accurately; however, the ultimate responsibility rests with the prescribing physician. Gastro-Hep Communications, Inc., the supporter, and the participants shall not be held responsible for errors or for any consequences arising from the use of information contained herein. Readers are strongly urged to consult any relevant primary literature. No claims or endorsements are made for any drug or compound at present under clinical investigation.

©2020 Gastro-Hep Communications, Inc., 611 Broadway, Suite 310, New York, NY 10012. Printed in the USA. All rights reserved, including the right of reproduction, in whole or in part, in any form.



# EDITORIAL ADVISORY BOARD

## EDITOR-IN-CHIEF:

**Gary R. Lichtenstein, MD**  
University of Pennsylvania

## SECTION EDITORS:

**Todd H. Baron, MD**  
University of North Carolina  
at Chapel Hill School of  
Medicine

**William D. Chey, MD**  
University of Michigan Medical  
Center

**Robert G. Gish, MD**  
Stanford University

**Stephen B. Hanauer, MD**  
Northwestern University Feinberg  
School of Medicine

**Stephen A. Harrison, MD**  
University of Oxford

**Eugene R. Schiff, MD**  
University of Miami Miller School  
of Medicine

**Prateek Sharma, MD**  
University of Kansas School of  
Medicine

---

**Maria T. Abreu, MD**  
University of Miami School of  
Medicine

**Leonard Baidoo, MD**  
Northwestern University  
Feinberg School of Medicine

**Robert N. Baldassano, MD**  
Children's Hospital of Philadelphia  
University of Pennsylvania

**Manoop S. Bhutani, MD**  
University of Texas  
M. D. Anderson Cancer Center

**Athos Bousvaros, MD, MPH**  
Children's Hospital Boston

**Joel V. Brill, MD**  
Predictive Health, LLC

**Robert S. Brown Jr, MD, MPH**  
Weill Cornell Medical College

**Brooks D. Cash, MD**  
University of Texas Health Science  
Center at Houston

**Lin Chang, MD**  
David Geffen School of Medicine  
University of California,  
Los Angeles

**Russell D. Cohen, MD**  
University of Chicago

**Scott J. Cotler, MD**  
University of Illinois at Chicago

**Douglas Dieterich, MD**  
Mount Sinai Medical Center

**Adrian M. Di Bisceglie, MD**  
Saint Louis University

**Jack A. Di Palma, MD**  
University of South Alabama

**David B. Doman, MD**  
George Washington University  
School of Medicine

**Herbert L. DuPont, MD**  
University of Texas McGovern  
Medical School  
University of Texas School of  
Public Health

**Gary W. Falk, MD**  
University of Pennsylvania Perelman  
School of Medicine

**Ronnie Fass, MD**  
Case Western Reserve  
University

**Brian G. Feagan, MD**  
Western University

**M. Brian Fennerty, MD**  
Oregon Health & Science  
University

**Steven L. Flamm, MD**  
Northwestern University  
Feinberg School of Medicine

**Basavana Goudra, MD**  
University of Pennsylvania

**Tarek Hassanein, MD**  
University of California,  
San Diego

**Colin W. Howden, MD**  
University of Tennessee Health  
Science Center

**Ira M. Jacobson, MD**  
Icahn School of Medicine at  
Mount Sinai

**David L. Jaffe, MD**  
University of Pennsylvania  
School of Medicine

**Lennox J. Jeffers, MD**  
University of Miami

**Maureen M. Jonas, MD**  
Boston Children's Hospital

**Sunanda V. Kane, MD, MSPH**  
Mayo Clinic

**Philip O. Katz, MD**  
Weill Cornell Medicine

**Seymour Katz, MD, FAGG,  
MACG**  
New York University

**Asher Kornbluth, MD**  
Mount Sinai Medical Center

**Joshua Korzenik, MD**  
Brigham and Women's  
Hospital

**Brian E. Lacy, MD, PhD**  
Mayo Clinic

**Anthony J. Lembo, MD**  
Beth Israel Deaconess Medical  
Center

**Richard MacDermott, MD**  
Albany Medical Center

**Willis C. Maddrey, MD**  
University of Texas Southwestern  
Medical Center

**Paul Martin, MD**  
University of Miami

**Kevin D. Mullen, MD**  
Metrohealth Medical Center

**Guy W. Neff, MD, MBA**  
Florida Research Institute

**Marion G. Peters, MD**  
University of California, San  
Francisco

**Mark Pimentel, MD, FRCP(C)**  
Cedars-Sinai Medical Center

**Paul J. Pockros, MD**  
Scripps Clinic

**Fred Poordad, MD**  
Texas Liver Institute/University of  
Texas Health, San Antonio

**Eamonn M. M. Quigley, MD**  
Houston Methodist  
Hospital

**K. Rajender Reddy, MD**  
University of Pennsylvania

**Douglas K. Rex, MD**  
Indiana University  
Medical Center

**Joel E. Richter, MD, FACP,  
MACG**  
University of South Florida

**David T. Rubin, MD**  
University of Chicago

**Paul Rutgeerts, MD**  
Katholieke Universiteit  
Leuven

**Sammy Saab, MD, MPH**  
David Geffen School  
of Medicine  
University of California,  
Los Angeles

**Seymour M. Sabesin, MD**  
Rush University  
Medical Center

**William J. Sandborn, MD**  
University of California  
San Diego

**Ellen J. Scherl, MD**  
Weill Cornell Medicine  
New York-Presbyterian  
Hospital

**Philip S. Schoenfeld, MD,  
MEd, MSc**  
John D. Dingell VA  
Medical Center

**Bo Shen, MD**  
The Cleveland Clinic

**Mitchell Shiffman, MD**  
Liver Institute of Virginia Bon  
Secours Health System

**Corey A. Siegel, MD**  
Dartmouth-Hitchcock Medical  
Center

**Jerome H. Siegel, MD**  
Mount Sinai Beth Israel

**Mark Sulkowski, MD**  
Johns Hopkins University  
School of Medicine

**Nicholas J. Talley, MD, PhD**  
Mayo Clinic

**Michael F. Vaezi, MD, PhD**  
Vanderbilt University  
Medical Center

**Fernando Velayos, MD**  
University of California,  
San Francisco

**Nizar Zein, MD**  
Cleveland Clinic  
Foundation

# INFORM UC: An Update on Contemporary Management of UC

## Dr Maria T. Abreu

Professor of Medicine  
Professor of Microbiology  
and Immunology  
Martin Kalser Chair in Gastroenterology  
Director, Crohn's & Colitis Center  
University of Miami  
Miller School of Medicine  
Miami, Florida

## Dr Stephen B. Hanauer

Clifford Joseph Barborika Professor  
of Medicine  
Medical Director, Digestive Health  
Center  
Northwestern University Feinberg  
School of Medicine  
Chicago, Illinois

## Dr Gary R. Lichtenstein


Director, Center for Inflammatory Bowel  
Disease  
The University of Pennsylvania Health  
System  
Professor of Medicine  
The Raymond and Ruth Perelman School of  
Medicine of the University of Pennsylvania  
Philadelphia, Pennsylvania

## Introduction

In July 2020, a group of clinical experts in the field of inflammatory bowel disease (IBD) convened virtually to discuss current issues and advances in the management of ulcerative colitis (UC). The faculty

began by reviewing abstracts from Digestive Disease Week (DDW) 2020 with particular relevance to the understanding and management of UC. Key advances in the contemporary management of UC were then reviewed, followed by an update on coronavirus disease 2019 (COVID-

19) and its implications for patients with IBD. This supplement, which summarizes discussions from the meeting, is intended to provide clinicians with the latest information regarding current treatment options and paradigms for UC.



**Figure 1.** Dietary patterns associated with increased risk of Crohn's disease. Adapted from Lo et al<sup>1</sup> and Nguyen et al.<sup>2</sup>

## UC Highlights From DDW 2020

### *Pathogenesis, Natural History, and Epidemiology for Clinicians*

#### Dietary Patterns and Risk of IBD

The effects of various dietary patterns on the risk of developing IBD were assessed in 2 studies using data from participants from the Nurses' Health Studies and Health Professionals Follow-Up Study.<sup>1,2</sup> Dietary information was obtained from validated questionnaires administered every 4 years, and self-reported diagnoses of IBD were confirmed through review of medical records. Lo and colleagues used this information to calculate an empirical dietary inflammatory pattern (EDIP) score, based on a weighted sum of 18 food groups (Figure 1).<sup>1</sup> In this analysis, participants with the most inflammatory diet (ie, highest quartile of cumulative average EDIP score) had a 45% increase in risk of Crohn's disease (CD) (hazard ratio, 1.45; 95% CI, 1.06-1.99) compared with those in the lowest quartile. In addition, dietary information was used to calculate a score for each participant's adherence to the sulfur microbial diet, an intake pattern associated with increased sulfur-metabolizing bacteria and characterized by intake high in processed meats and low in mixed vegetables and legumes (Figure 1).<sup>2</sup> As with the inflammatory diet, participants in the top quartile of sulfur microbial diet scores had an increased risk of CD compared with participants in the lowest quartiles, with a pooled multivariate relative risk of 1.49 (95% CI, 1.07-2.08; *P* trend=.01). Neither the inflammatory diet nor the sulfur microbial diet was significantly associated with the risk of developing subsequent UC.<sup>1,2</sup> While the cohort of nurses in these studies only reflects women and is somewhat older, and less heterogeneous than the US population of IBD patients, these data suggest the possibility of proinflammatory food groups contributing to sulfur-metabolizing bacteria and the risk of CD.

**Table 1.** Risk of Adverse Outcomes in Patients With Any 1 of 3 Markers of Low Social Status During the Study Period

	HR (95% CI)
Hospitalizations	1.54 (1.45-1.64)
ICU admission	1.90 (1.62-2.23)
Mortality	1.49 (1.33-1.69)
High-dose corticosteroid	1.20 <sup>a</sup> (1.11-1.29)

<sup>a</sup>Relative risk. HR, hazard ratio; ICU, intensive care unit. Adapted from Bernstein et al.<sup>3</sup>

#### Social Determinants of Health in IBD


Consistent with increasing evidence of health disparities in IBD, analysis of data from patients with IBD in Manitoba, Canada demonstrated a significant association between markers of low social status and poor outcomes in IBD.<sup>3</sup> In this study, 2905 (31.2%) of 9298 Manitoba residents with IBD were found to have at least 1 indicator of low social status (income assistance, children with endangered health or emotional well-being, and/or composite score reflecting household income and education rates). Having any of these 3 markers of low social status was linked to higher risk of hospitalization, intensive care unit (ICU) admission, high-dose corticosteroid use, and a higher rate of mortality (Table 1). These findings are consistent with other studies suggesting that socioeconomic factors contribute to poor outcomes in IBD, including hospitalizations and high corticosteroid use.<sup>4-6</sup>

#### Microbial Dysbiosis in Pregnant Women and Their Infants

An analysis of data from the MECONIUM (Exploring Mechanisms of Disease Transmission in Utero Through the Microbiome) study highlighted potential effects of the altered microbiome on IBD activity in pregnant women and on their offspring during the first few years of life.<sup>7</sup> Fecal calprotectin levels of 341 pregnant women (90 with IBD, 251 controls) were measured at each trimester of pregnancy and in their babies (n=290) throughout the first 3 years of life. Fecal calprotectin levels were significantly higher among mothers with IBD (*P*<.001 at every trimester) than control mothers and declined during pregnancy. Higher fecal calprotectin in the babies was associated with maternal IBD (particularly with active disease), lower microbiome diversity, and the abundance of certain microbial genera (*Blautia* and *Streptococcus*). These data from the MECONIUM study suggest not only that alterations in the microbiome may affect IBD activity during pregnancy, but also that infants born of mothers with IBD, especially active IBD, demonstrate signs of intestinal inflammation in the neonatal period.

#### Genetic Associations and EIMs in IBD

Genotype data from 3 large databases were analyzed in an effort to better understand the pathogenesis of extraintestinal manifestations (EIMs) in IBD.<sup>8</sup> Mega case-control regression analyses were conducted to compare cases with at least 1 joint, skin, eye, or




**Figure 2.** Clinical remission among week 8 responders at week 52 (primary endpoint) in the SERENE-UC study. EOW, every other week; SERENE-UC, Study to Evaluate the Safety and Efficacy of Two Drug Regimens in Subjects With Moderate to Severe Ulcerative Colitis; TDM, therapeutic drug monitoring. Adapted from Colombel et al.<sup>9</sup>

liver EIM (n=884) and those without EIMs (n=11,029). Several EIM associations were identified at established IBD susceptibility loci as well as novel loci. Further, logistic regression of classical human leukocyte antigen (HLA) alleles demonstrated significant associations of ankylosing spondylitis at the known alleles HLA-B27 and HLA-C\*02. Psoriasis was associated with the known HLA-C\*06 allele as well as 2 novel HLA alleles. These findings highlight the power of pooling genetic data to identify genetic pathways linked to EIMs in IBD, although better phenotyping of the patients with respect to EIMs will be needed.

### Clinical and Therapeutic Advances

**Optimizing Current Therapies** The SERENE-UC (Study to Evaluate the Safety and Efficacy of Two Drug Regimens in Subjects with Moderate to Severe Ulcerative Colitis) study was a phase 3, double-blind trial comparing higher vs standard adalimumab dosing regimens for induction and maintenance therapy in adults with moderate to severely active UC. After completion of the 8-week induction study, 757 patients were rerandomized to receive adalimumab 40 mg every week, every other week, or using therapeutic drug monitoring (TDM) regimens.<sup>9</sup> Analysis of the primary endpoint in the intent-to-treat responder population (n=371) demonstrated numerically, but not significantly, higher rates of clinical remission (Full Mayo Score  $\leq 2$  with no subscore  $>1$ ) among patients receiving adalimumab every week than every other week (39.5% vs 29.0%, respectively;  $P=.069$ ) (Figure 2). Weekly maintenance dosing was associated with higher rates of corticosteroid-free patients than every-other-week dosing (74.7% vs 53.3%;  $P=.002$ ), but other secondary efficacy endpoints were similar between groups. The safety profile was similar between treatment arms, and no new long-term safety concerns were identified. Overall, these data reinforce the efficacy of the current US Food and Drug Administration–approved




**Figure 3.** Rates of ADAs by anti-TNF exposure. ADA, antidrug antibody; TNF, tumor necrosis factor. Adapted from Yanai et al.<sup>11</sup>

maintenance regimen for adalimumab in UC<sup>10</sup> and demonstrate that more intensive maintenance therapy does not significantly improve outcomes over time. However, it is important to remember that controlled trials such as this do not allow personalized dose adjustments that are frequently needed in clinical practice based on either reactive or proactive TDM.

Given the practical and safety implications of immunogenicity in patients treated with anti-tumor necrosis factor (TNF) agents, Yanai and colleagues examined the rate of consecutive immunogenicity to a second anti-TNF relative to antidrug antibody (ADA) formation against the first agent.<sup>11</sup> In this retrospective cross-sectional study, the immunogenicity rates of 55 patients from 2 US medical centers who switched from 1 anti-TNF agent to another were analyzed and compared with those of the overall ADA rates determined from 1570 tests in the laboratory database. Of 25 patients who developed ADAs to infliximab as their first anti-TNF agent, 7 (28%) developed ADAs against adalimumab after switching to this agent. Seven of 10 (70%) patients with ADAs against adalimumab developed ADAs to infliximab after switching. Regardless of which anti-TNF agent was used first, the rates of consecutive immunogenicity were significantly higher than the overall immunogenicity rates (28% consecutive adalimumab immunogenicity vs 13% overall adalimumab immunoge-

nicity;  $P=.03$ ; 70% consecutive infliximab immunogenicity vs 29% overall infliximab immunogenicity;  $P=.007$ ) (Figure 3). Only 33% of patients were treated with concomitant immunomodulators after switching anti-TNF therapies due to immunogenicity. These findings suggest that patients with consecutive immunogenicity may be predisposed to developing ADAs to anti-TNF therapies, and underscore the need for concomitant immunomodulator therapy in such patients.

Recognizing the need for real-world data, the effects of tofacitinib on endoscopic and histologic outcomes in patients were prospectively evaluated in a cohort of 35 patients with UC refractory to anti-TNF agents and vedolizumab.<sup>12</sup> After treatment with tofacitinib 10 mg twice daily for 8 weeks, the Mayo endoscopic subscore decreased significantly from baseline ( $P=.004$ ). At week 8, 22.9% of patients achieved endoscopic response (Mayo endoscopic subscore  $\leq 1$ ),



**Figure 4.** Week 48 remission among patients with baseline Mayo endoscopic subscore of 0 or 1. Adapted from Kobayashi et al.<sup>14</sup>

17.2% achieved endoscopic remission (Mayo endoscopic subscore of 0), and 14.8% achieved histologic remission (numeric Geboes score  $\leq 6$ ). Biologic response (50% decrease in fecal calprotectin or fecal calprotectin  $< 250$  mg/g) and remission (fecal calprotectin  $< 250$  mg/g at week 8) were achieved in 52.9% and 38.2% of patients, respectively. Multivariate analysis found that higher baseline serum albumin levels and lower Mayo endoscopic subscores were independent predictors of endoscopic and biologic remission. Primary nonresponse to tofacitinib was observed in 10 of 20 (50%) patients with primary nonresponse to 1 anti-TNF agent, 4 of 5 (80%) patients with primary nonresponse to 2 anti-TNF agents, and 3 of 8 (37.5%) patients with primary nonresponse to vedolizumab. These data support the efficacy of tofacitinib in inducing biologic, endoscopic, and histologic remission in patients with refractory UC, although it does not appear to be an alternative treatment strategy for those with primary nonresponse to 2 anti-TNF agents. Commenting on these data, Dr Stephen B. Hanauer noted that “given that tofacitinib is more effective in the anti-TNF-naïve populations than those with previous exposure, it is not surprising that in this small series of patients, more refractory patients (based on prior exposure to more than 1 biologic) responded less well than biologic-naïve patients.”

The impact of combination therapy with immunomodulators (azathioprine, 6-mercaptopurine, and methotrexate) was retrospectively evaluated in a cohort of 549 patients (263 with UC, 286 with CD) receiving maintenance therapy with vedolizumab and 363 patients (4 with UC, 359 with CD) receiving ustekinumab maintenance therapy across 3 tertiary centers.<sup>13</sup> Of these, 131 patients receiving vedolizumab and 120 receiving ustekinumab were receiving combination therapy with thiopurines or methotrexate. At week 14, combination therapy did not achieve superior clinical response or remission

“This study shows that patients who have a planned withdrawal of the biologic have a 50% chance of flaring. It is my hope that a combination of microbiome and microbiome metabolism analyses could predict whether a patient will stay in a long-lived remission, even after stopping a biologic or a small molecule.”

– Dr Maria T. Abreu

rates (primary endpoint) compared with either vedolizumab monotherapy (68.2% vs 74.1%, respectively;  $P=.22$ ) or ustekinumab monotherapy (54.6% vs 65.8%;  $P=.08$ ). Similarly, clinical response or remission rates did not differ between treatment groups at week 30 or 54. Combination therapy was not found to improve endoscopic response at 1 year compared with

monotherapy. Although these results fail to support a benefit of combining immunomodulators with vedolizumab or ustekinumab, prospective, properly powered studies are needed to define the role of combination therapy in patients treated with non-anti-TNF biologics. Dr Hanauer commented that these results “add to the accumulating real-world and clinical trial data suggesting that there is less impact on outcomes with combination therapy vs monotherapy with ustekinumab and vedolizumab, both of which are inherently less immunogenic than infliximab and adalimumab.”

**Predicting UC Relapse** In the first prospective, randomized trial evaluating the impact of infliximab discontinuation in quiescent UC, 92 patients with UC maintained in clinical remission with infliximab were randomized to continue or discontinue infliximab in a 1:1 ratio.<sup>14</sup> Eligible patients were confirmed to be in clinical remission for longer than 6 months, corticosteroid-free, and had Mayo endoscopic subscores of 0 or 1. At week 48, remission rates were significantly higher in patients who continued infliximab compared with those who discontinued treatment (80.4% vs 54.3%, respectively;  $P=.008$ ) (Figure 4). While the Nancy histologic index and C-

**Table 2.** Adjusted HRs of Pneumonia and Cox Model Results Adjusted for All Covariates

	Adjusted HR (95% CI)	P value
Charlson Comorbidity Index	1.16 (1.14-1.18)	<.001
Influenza vaccination	1.28 (1.19-1.38)	<.001
Narcotic within 60 days prior to index date	1.45 (1.34-1.57)	<.001
Prednisone cumulative (mg/day)	1.02 (1.01-1.03)	<.001
Prednisone within 30 days prior to index date	1.99 (1.78-2.22)	<.001
IBD flare	2.64 (2.38-2.93)	<.001
PPSV23	1.15 (1.07-1.23)	<.001
Medications (5-ASA reference)		
Thiopurines	0.92 (0.81-1.04)	.173
Anti-TNF	1.21 (1.05-1.40)	.01
Thiopurine/anti-TNF combination	1.12 (0.84-1.51)	.439
Vedolizumab	0.74 (0.30-1.80)	.508

5-ASA, 5-aminosalicylate; HR, hazard ratio; IBD, inflammatory bowel disease; PPSV23, pneumococcal polysaccharide vaccine; TNF, tumor necrosis factor. Adapted from Patel et al.<sup>20</sup>

reactive protein levels were predictive of remission at week 48, use of concomitant immunomodulators, serum infliximab trough concentrations, and Mayo endoscopic subscore of 0 at baseline were not. Most (66.7%) of the patients who relapsed achieved remission within 8 weeks of infliximab retreatment. These findings confirm that discontinuation of maintenance infliximab increases the risk of relapse in UC and suggest that endoscopic normalization does not guarantee successful infliximab discontinuation.

The value of various protein, metabolite, or microbial markers in predicting IBD relapse was prospectively explored in a cohort of 164 patients with quiescent IBD (108 with CD, 56 with UC).<sup>15</sup> Quiescent disease was defined as absence of clinical symptoms (Harvey-Bradshaw Index <4, Simple Clinical Colitis Activity Index <2) and endoscopic remission assessed by colonoscopy within the previous year. Serum samples were used to perform metabolomic and proteomic profiling, and stool samples were used for metagenomic sequencing. Multivariate models found 3 protein biomarkers (interleukin

[IL]-10, GDNF, and CD8A) and 4 metabolic markers (propionyl-L-carnitine, carnitine, sarcosine, and sorbitol) to be significantly associated ( $P<.05$ ) with relapse (symptomatic worsening accompanied by elevated inflammatory markers resulting in a change in therapy or IBD-related hospitalization or surgery). High proteomic (odds ratio [OR], 9.11; 95% CI, 1.90-43.61) and metabolomic (OR, 5.79; 95% CI, 1.24-27.11) risk scores independently predicted higher risk of relapse over 2 years. Further, proteomic and metabolomic changes associated with relapse correlated with proinflammatory changes in the microbiome.

**Safety of IBD Therapies** A number of studies reported at DDW 2020 add to our knowledge of the safety of current IBD therapies. The risk of lymphoma was evaluated among a population-based cohort of IBD patients in Israel (N=37,549), 164 of whom had been diagnosed with lymphoma.<sup>16</sup> Multivariate analysis demonstrated that older age at follow-up start ( $P=1.3 \times 10^{-8}$ ) and recent (up to 6 months) thiopurine exposure

( $P=.004$ ) were associated with incident lymphoma diagnosis, whereas ever-exposure to thiopurines alone or in combination were not. The association of lymphoma with recent thiopurine exposure is notable given previous studies reporting an increased risk of lymphoma with increasing thiopurine exposure.<sup>17,18</sup> Further, the findings from this cohort differ from previous research that has demonstrated higher lymphoma risk with combination thiopurine and anti-TNF therapy compared with either monotherapy.<sup>19</sup>


The impact of immunosuppressive therapies on the risk of pneumonia was examined in a large nationwide cohort of IBD patients in the Veteran Affairs dataset from 2000 through 2019.<sup>20</sup> Of 56,398 patients with IBD in the database, 3442 (6.1%) developed all-cause pneumonia during a median 8 years of follow-up. Multivariate Cox regression analysis demonstrated an increased risk of pneumonia with exposure to anti-TNF therapy compared with 5-aminosalicylate (5-ASA) alone, higher cumulative corticosteroid dose, and any corticosteroid use in the preceding 30 days (Table 2). In contrast, thiopurines, vedolizumab, or combina-

**Table 3.** DVT and PE IRs Among Patients in the Tofacitinib Clinical Trial Program

	Induction Cohort (8 weeks) <sup>a</sup>		Maintenance Cohort (52 weeks) <sup>a</sup>			Overall Cohort (≤6.8 years) <sup>a,b</sup>		
	Placebo	Tofacitinib 10 mg BID	Placebo	Tofacitinib 5 mg BID	Tofacitinib 10 mg BID	PD Tofacitinib 5 mg BID	PD Tofacitinib 10 mg BID	Tofacitinib All
N	282	938	198	198	196	198	959	1157
PY	44.8	156.2	100.4	146.2	154.3	664.1	1917.1	2581.3
DVT								
n (%)	1 (0.4)	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.1)	1 (0.1)
IR	1.99	0.00	0.97	0.00	0.00	0.00	0.05	0.04
(95% CI)	(0.1-11.1)	(0.0-2.2)	(0.0-5.4)	(0.0-2.5)	(0.0-2.4)	(0.0-0.5)	(0.0-0.3)	(0.0-0.21)
PE								
n (%)	1 (0.4)	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)	0 (0.0)	4 (0.4)	4 (0.3)
IR	1.98	0.00	0.98	0.00	0.00	0.00	0.20	0.15
(95% CI)	(0.1-11.0)	(0.0-2.2)	(0.0-5.4)	(0.0-2.5)	(0.0-2.4)	(0.0-0.5)	(0.1-0.5)	(0.0-0.4)

For Overall Cohort analysis, patients were categorized based on the average daily dose of tofacitinib (placebo exposure was not included): PD tofacitinib 5 mg BID (average total daily dose <15 mg) and PD tofacitinib 10 mg BID (average total daily dose ≥15 mg). <sup>a</sup>Events occurring up to 28 days beyond the last dose of the corresponding cohort are included in the analyses. <sup>b</sup>Data as of May 2019 (OLE study database not locked). BID, twice daily; DVT, deep vein thrombosis; IR, incidence rate; N, number of patients; n, unique number of patients with a particular adverse event; OLE, open-label, long-term extension; PD, predominant dose; PE, pulmonary embolism; PY, patient-years. Adapted from Sandborn et al.<sup>22</sup>





**Figure 5.** Mean trough (predose) infliximab concentrations with CT-P13 SC or IV. IV, intravenous; SC, subcutaneous. Adapted from Reinisch et al.<sup>26</sup>

tion thiopurine and anti-TNF therapy were not significantly associated with a greater risk of pneumonia. Nonmedication-related factors associated with increased pneumonia risk included older age, male sex, more comorbidities, IBD flares, and higher health care utilization.


In light of recent updates to the tofacitinib prescribing information regarding the risk of thromboembolism,<sup>21</sup> Sandborn and colleagues reported an update on the incidence of deep vein thrombosis (DVT) and pulmonary embolism (PE) in the tofacitinib UC clinical development program.<sup>22</sup> A total of 1157 patients involved in placebo-controlled induction/maintenance studies and/or an open-label, long-term extension (OLE) study were evaluated for DVT and PE, with 2581.3 patient-years of tofacitinib exposure and up to 6.8 years of treatment. Four patients in the Induction and Maintenance Cohorts experienced a DVT or PE, and all were receiving placebo at the time of the event (Table 3). Five patients, all receiving tofacitinib 10 mg twice daily, in the Overall Cohort experienced a thromboembolic event during the OLE study. These events occurred after 7 months or more of treatment, and all patients had risk factors (other than UC) for venous thrombosis. These results indicate that the incidence rates

of thromboembolic events in the Overall Cohort have remained stable since last reported.<sup>23</sup>

**Looking to the Future** Recognizing the therapeutic potential of low doses of IL-2 in immune-mediated diseases as well as in preclinical models of UC,<sup>24</sup> a phase 1b/2a study was conducted to examine the safety and efficacy of this agent when administered subcutaneously to patients with moderate to severely active UC.<sup>25</sup> In this open-label, single-arm trial, 24 patients with UC and a Mayo score of 6 to 12 received daily subcutaneous (SC) IL-2 during a dose-escalation phase followed by treatment with the maximum tolerated dose. The most common (occurring in >10% of patients) adverse events across all doses included injection site reactions and malaise, and no serious adverse events were observed. Ten of 24 (41.6%) patients achieved either response or remission. A dose of  $1 \times 10^6$  IU/m<sup>2</sup>/day was found to be the maximum effective dose, achieving a biologic response and peripheral regulatory T-cell expansion without significant toxicity.

CT-P13 SC, a novel infliximab formulation approved by the European Medicines Agency for use in rheumatoid arthritis, has demonstrated comparable clinical efficacy

and safety with CT-P13 intravenous (IV) in a controlled trial involving patients with CD and UC for up to 30 weeks.<sup>26</sup> Reinisch and colleagues reported further data from this trial, characterizing the efficacy, pharmacokinetics, and safety of CT-P13 SC over 1 year. After receiving 5 mg/kg IV loading doses at weeks 0 and 2, patients were randomized at week 6 to either SC 120 mg (patients <80 kg) or 240 mg (patients ≥80 kg) every 2 weeks or to continue 5 mg/kg IV every 8 weeks. At week 30, patients in the IV arm were switched to the SC arm. A total of 105 of 131 randomized patients completed the week 54 visit. Clinical response and remission rates were comparable between the IV and SC arms at week 30 and at week 54 after the remaining IV patients switched to SC. Rates of mucosal healing were also similar between arms at week 54. The overall safety profiles of the formulations were comparable, with the exception of a higher rate of localized injection site reactions in the SC group. Trough (predose) infliximab concentrations in the IV arm increased to levels similar to those in the SC group and were maintained through week 54 (Figure 5). These results support CT-P13 as a viable therapeutic option for use early after induction or in patients switching from IV during maintenance therapy.



## Update on UC Management

The recognition of the importance of minimizing disease activity early in the disease course, coupled with an increasing ability to measure clinical and objective endpoints, has led to evolving treatment paradigms in UC. A key emerging strategy is a treat-to-target approach, which aims to achieve disease remission by adjusting therapy according to the achievement of treatment response targets.<sup>27-29</sup> This concept has widespread adaptation in other diseases, such as targeting normal blood pressure in patients with hypertension or normal glycosylated hemoglobin in patients with diabetes. From a practical perspective, the treat-to-target strategy requires collaboration between the physician and patient to assess baseline disease characteristics, identify appropriate targets, tailor therapy based on the risk of disease progression, monitor progress, and optimize therapy to reach the goal (Figure 6).<sup>27</sup>

### Assessing Disease Severity and Activity

Once a diagnosis of UC is made, assessment of both disease activity and severity are essential in formulating a treatment plan.<sup>30</sup> In differentiating

**Figure 6.** Proposed ulcerative colitis treat-to-target algorithm. <sup>a</sup>Mucosal healing as a treatment target must involve patient decision because of the high burden of monitoring and potential need for therapy escalation despite symptom resolution. <sup>b</sup>Biomarker normalization as a treatment target must involve patient decision because of potential need for therapy escalation despite symptom resolution. <sup>c</sup>If adjunctive biomarkers are not improving or normalizing, mucosal healing targets should be reassessed. RHI, Roberts Histopathology Index; UCEIS, Ulcerative Colitis Endoscopic Index of Severity. Adapted from Ungaro et al.<sup>29</sup>

**Table 4.** American College of Gastroenterology Guidelines for Severity in Active Ulcerative Colitis

	Remission	Mild	Moderate to Severe	Fulminant
Stools (number/day)	Formed stools	<4	>6	>10
Blood in stools	None	Intermittent	Frequent	Continuous
Urgency	None	Mild, occasional	Often	Continuous
Hemoglobin	Normal	Normal	<75% of normal	Transfusion required
ESR (mm/hr)	<30	<30	>30	>30
CRP (mg/L)	Normal	Elevated	Elevated	Elevated
FC (µg/g)	<150-200	>150-200	>150-200	>150-200
Endoscopy (Mayo subscore)	0-1	1	2-3	3
UCEIS	0-1	2-4	5-8	7-8

The above factors are general guides for disease activity. With the exception of remission, a patient does not need to have all of the factors to be considered in a specific category. CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; FC, fecal calprotectin; UCEIS, Ulcerative Colitis Endoscopic Index of Severity. Adapted from Rubin et al.<sup>30</sup>

“A treat-to-target strategy is not a snapshot in the time course of a disease but rather an iterative, ongoing process of assessing the parameter, modifying treatment, subsequent assessment of response, and further modifications until the endpoint is achieved and then maintained. The process does not stop when the target is met but is ongoing to ensure long-term stability.”

– Dr Stephen B. Hanauer

these 2 parameters, Dr Hanauer explained that “disease activity can be considered a snapshot in time, capturing clinical signs and symptoms that are independent of ongoing medications or surgeries.” Whereas activity does not imply the clinical course, disease severity considers the longitudinal factors associated with the prognosis. Based on the most recent American College of Gastroenterology guidelines on UC management, disease severity in active disease ranges from mild to severe based on a combination of patient-reported outcomes, laboratory values, and endoscopic parameters (Table 4). However, these factors actually reflect disease activity rather than prognosis, aside from the presence of deep ulcers reflected by a Mayo 3 endoscopic subscore. Thus, disease severity reflects not only symptoms, but factors that contribute to the risk of disease progression and influence response to various therapies.

### *Treatment Targets*

Recognizing the disabling nature of UC, treatment goals are increasingly focused on slowing disease progression with the hope of altering the natural history of the condition.<sup>28-30</sup> To that end, treating to symptom resolution alone may leave active disease (ie, mucosal inflammation) and has

“When practitioners choose therapies for patients with UC, it is important to take into consideration not only their disease activity (that is, how sick they are at the time they have active disease), but their prognosis. A practitioner may see a patient with mild activity but a poor prognosis and start that patient on initial therapy with biologic therapy. The recent ACG guidelines for the first time also take into consideration the presence of fecal urgency. In addition, fecal calprotectin and endoscopic disease activity are included. Thus, a conglomerate of endoscopy, symptoms, and biomarkers may help a practitioner decide on appropriate therapeutic interventions.”

– Dr Gary R. Lichtenstein

not been proven sufficient to alter long-term remission or complication rates.<sup>29</sup> To the contrary, a proportion of patients with UC has been shown to have mucosal inflammation without clinical symptoms.<sup>30,31</sup> “With the introduction of endoscopic capabilities and indices that reproduce and validate endoscopic findings in the colon,” Dr Hanauer explained, “discrepancies between the severity of symptoms, objective markers of inflammation, and the endoscopic findings became apparent.” Moreover, a number of studies have demonstrated that achieving mucosal healing, or endoscopic remission, is associated with improved outcomes in UC, including lower risk of colectomy, lower clinical relapse and hospitalization rates, and reduced rates of dysplasia and colorectal cancer.<sup>32,33</sup> In light of these findings, treatment goals have evolved beyond symptom resolution alone to include sustained control of inflammation, with mucosal healing now widely recognized as an important goal of therapy.<sup>28-30</sup>

Although the endoscopic aspects of mucosal healing have emerged as a primary treatment goal in UC, the histologic aspects of mucosal healing are just beginning to be appreciated.<sup>29,30,34-37</sup> Histologic healing, defined as microscopic normalization of the colonic mucosa, is a distinct target from endoscopic healing, a measure of endoscopically visible disease activity.<sup>29</sup> Dr Hanauer noted that “with validation of histologic scores in UC, such as the Robarts or Nancy histologic indices and Geboes score, resolution of histologic inflammation has been associated with additional improvements in outcomes.” Indeed, a growing number of studies demonstrate that histologic remission in UC is predictive of corticosteroid-free remission, clinical relapse, hospitalization, and corticosteroid use,<sup>30,38</sup> while others have correlated the degree of histologic inflammation with dysplasia and colorectal cancer.<sup>30,39,40</sup> Although more data are needed to identify which histologic indices should be used in research and in clinical practice,

Dr Maria T. Abreu noted that “mucosal healing, which includes both endoscopic and histologic normalization, should be an aspirational goal in treating our patients.”

In speaking to the clinical utility of these endpoints, Dr Gary R. Lichtenstein noted that “perfection may be the enemy of excellence.” “For example,” he continued, “a patient with a previous Mayo score of 4 who is now a Mayo 1 with persistent histologic inflammation and is asymptomatic does not represent a failure of medical therapy. In other words, not all patients evaluated are mandated to have a Mayo 0 and absent histologic inflammation as their treatment endpoint.” All endpoints need to be considered in the context of where patients were prior to therapy and which therapies have been used. Dr Lichtenstein concluded, “Thus, although the presence of persistent histologic inflammation in the aforementioned patient may confer a higher risk of disease flare in the subsequent 6 to 12 months, the treatment endpoint achieved in this patient may be considered to be appropriate and the outcome successful.”

The clinical utility of adjunctive biomarkers, particularly fecal calprotectin, as targets in UC has grown, with accumulating data demonstrating correlations of low concentrations with the absence of mucosal inflammation and rising concentrations with relapse.<sup>29,30</sup> In contrast, advanced imaging modalities (eg, computed tomography, magnetic resonance, intestinal ultrasound) are useful in staging disease location and evaluating complications, but have not been adequately evaluated or established as surrogate targets for disease severity or response to therapies.<sup>29</sup> Dr Lichtenstein noted that although normalization of biomarkers is desirable, it is not necessarily the final endpoint to achieve, as verification with endoscopy is important. Normal C-reactive protein levels can be seen in patients with active disease<sup>41</sup> as well as in asymptomatic patients with mild

“Putting these all together has evolved the concept of a deep remission that includes the resolution of symptoms, endoscopic lesions, and biomarkers. Increasing numbers of clinical trials and real-world observational series have continued to support the concept that ‘the deeper the remission, the better the long-term course.’”

– Dr Stephen B. Hanauer

mucosal lesions, especially those with isolated involvement of the ileum.<sup>42</sup>

#### *Personalizing Therapy*

With the growing effort to slow disease progression, management of UC is increasingly driven by assessing a patient’s prognostic factors for aggressive disease.<sup>43-45</sup> With this approach, patients with risk factors for an unfavorable disease course are treated more aggressively after initial diagnosis than patients with fewer risk factors for progression, who may be managed with a conventional step-up approach.<sup>43,45</sup> In patients with limited anatomic involvement and mild endoscopic disease who are believed to have a low risk of colectomy, treatment with oral and/or rectal 5-ASAs with or without oral budesonide is recommended.<sup>30,46</sup> In contrast, more aggressive therapies are recommended for patients with poor prognostic factors such as extensive colitis, deep ulcers, previous requirement for corticosteroids, and failure to respond to conventional treatments. Options for such patients include initiating a thiopurine with a short

course of corticosteroids, a biologic agent (anti-TNF agent [infliximab, adalimumab, certolizumab pegol], vedolizumab, or ustekinumab) with or without an immunomodulator, or the oral Janus kinase (JAK) inhibitor tofacitinib.<sup>30,46,47</sup>

With the availability of several classes of biologics and targeted therapies with variable efficacy and safety profiles, positioning different agents in the treatment course of moderate to severe UC can be challenging.<sup>48</sup> Robust data support the efficacy of the anti-TNF therapies in achieving and maintaining clinical and endoscopic remission, improving quality of life, and reducing hospitalizations and surgeries in UC.<sup>30,49-54</sup> The most important safety concern with these agents is the risk of serious infection, which may be reduced by screening for hepatitis B and tuberculosis and ensuring appropriate immunization before initiating treatment.<sup>55</sup> Vedolizumab, a gut-selective agent that

“Perhaps as a result of the age of our patient population, and the fact that tofacitinib is being used primarily in the outpatient setting, the increase in thromboembolic complications seen in rheumatoid arthritis patients has not been seen in UC. This concern has limited the use of tofacitinib in the right population, which would be younger patients who could achieve rapid remission with this agent.”

– Dr Maria T. Abreu

acts by binding the  $\alpha_4\beta_7$  integrin,<sup>56,57</sup> has emerged as a first-line agent in this setting due to its efficacy, favorable safety profile, and low rate of immunogenicity.<sup>55,58</sup> Ustekinumab, an anti-p40 antibody that inhibits IL-12 and -23, induces and maintains remission in UC and, like vedolizumab, offers excellent safety with low immunogenicity.<sup>55</sup>

Unlike the biologics, tofacitinib is an oral small molecule inhibitor that has demonstrated a notably rapid onset of action, with some patients in phase 3 trials achieving significant improvement in the partial Mayo score as early as 2 weeks.<sup>59</sup> Dr Lichtenstein emphasized the importance of onset when choosing therapies and added that post hoc analysis of tofacitinib clinical trial data demonstrated benefit within 3 days of use. Given this rapid onset, patients hospitalized with acute severe UC have been successfully treated with tofacitinib.<sup>60</sup> “Although this use in hospitalized patients is currently an off-label use,” he noted, “future validation of clinical treatment efficacy in this study population might be of interest to assess in a prospective randomized clinical trial.” Key safety concerns with tofacitinib include the risk of infection, particularly reactivation of herpes zoster, hyperlipidemia, and thromboembolic risks.<sup>21,30,59</sup> Data indicating an increased risk of DVT and PE associated with tofacitinib in patients with rheumatoid arthritis recently prompted the addition of a warning to the product labeling.<sup>21</sup> Dr Abreu noted that tofacitinib has been relegated to second- or third-line use due to this thromboembolic risk, an unfortunate positioning “for many patients who could benefit from its mechanism of action and rapid response.”

The role of combination therapy in UC continues to be refined. Historically, clinical trials of biologics and the more recent JAK inhibitors enrolled patients with inadequate responses to aminosalicylates, immunosuppressives, or anti-TNF agents. Dr Hanauer noted that “within the context of these

trials where patients were refractory to their baseline medications, post hoc analyses did not demonstrate clinical benefits whether or not patients were receiving aminosalicylates or thiopurines (despite higher biologic drug levels and reduced ADAs).” He further explained that these trials were neither powered nor randomized according to baseline therapies. In contrast, subsequent data from trials such as the UC-SUCCESS study that randomized immunosuppressive- and bio-naïve patients clearly demonstrated a benefit of combining thiopurines and anti-TNF agents in UC.<sup>61</sup> Commenting on these findings, Dr Abreu noted that thiopurines not only increase the level of these biologics, but provide a complementary mechanism of action to TNF inhibition. Of interest, recent American Gastroenterological Association guidelines on managing moderate to severe UC suggest that vedolizumab or ustekinumab can also benefit from combination with immunosuppressives, a recommendation based on post hoc analyses.<sup>47</sup> Given that prospective, controlled trials have yet to be performed with non-TNF biologics, this recommendation has not been met with universal agreement.

### **Monitoring Progress**

Regular assessment of disease activity is an essential component of a treat-to-target strategy, although the frequency of monitoring varies by parameter and by disease activity and severity.<sup>28-30</sup> According to the STRIDE consensus statement, endpoints should be assessed at least every 3 months in patients with active disease and 6 to 12 months in patients who have achieved remission.<sup>28</sup> Symptoms are the most responsive to change and can be evaluated within days to weeks. Biomarkers such as C-reactive protein and fecal calprotectin can be assessed both short and long term. Expanding on this, Dr Lichtenstein noted that “laboratory evaluation every 4 weeks or less may be needed in patients with an acute flare, while assessments every 12 weeks may

be acceptable after patients have normalized after a recent flare or change in therapy.” In contrast, endoscopy and histology are longer-term targets that may require many months to demonstrate improvement. In general, patients starting on therapy for UC may generally have endoscopic confirmation of their target after 3 to 6 months,<sup>28</sup> perhaps with an intermediate biomarker. Dr Abreu added that “there is a role for colonoscopy for any major changes in medical therapy, and certainly at baseline to get the extent of disease, including biopsies of inflamed and normal-appearing mucosa.”

### **Optimizing Therapy**

With the evolution of validated treatment targets and expanded treatment options, the concept of optimizing therapies based on their pharmacokinetic and pharmacodynamic properties has become an important tool in managing UC. However, as Dr Hanauer noted, “this is not a novel concept in medicine or in IBD. Remember that weight-based dosing of thiopurines has been shown to be inferior to thioguanine levels in improving outcomes and reducing complications, and serum cyclosporine concentrations are useful in monitoring for both efficacy and safety.” More recently, however, growing evidence linking high concentrations of biologic agents with favorable outcomes (clinical, biomarker, and endoscopic remission),<sup>62-65</sup> as well as ADAs to worse outcomes (lower serum drug concentrations, reduced clinical response, and infusion reactions),<sup>66,67</sup> has established TDM as an important tool in optimizing UC management. Although much of these data have been obtained in patients on anti-TNF therapies,<sup>62-65</sup> exposure-response relationships have been demonstrated for other biologics, including vedolizumab<sup>68,69</sup> and ustekinumab.<sup>70</sup>

TDM can be used at any point in induction or maintenance therapy, either in a proactive routine fashion when the patient is in remission, or as a reactive strategy to help guide treatment in cases of suboptimal

response.<sup>71</sup> Although most current evidence has explored reactive TDM, recent retrospective data suggest that proactive monitoring of serum infliximab concentrations may be associated with better clinical outcomes and less need for IBD-related surgery or hospitalization compared with reactive monitoring.<sup>72</sup> In contrast, the TAXIT (Trough Concentration Adapted Infliximab Treatment) study in patients with CD found that, after initial infliximab dose optimization, the proportion of patients achieving remission at 1 year did not differ between those who received routine proactive TDM and those who received no TDM.<sup>73</sup> Despite these inconsistencies, Dr Abreu noted that proactive monitoring allows for necessary dosing adjustments to be made early in the course of treatment.

Another important aspect of optimizing UC management is to continually reevaluate the patient's medication regimen, adjusting dosing as necessary. For example, Dr Abreu noted, "there are patients who are finally doing well after a severe flare in whom the biologic dose or frequency can be reduced." The decision to discontinue a therapy, however, may be unclear, as definitive data to guide such decisions are lacking.<sup>74-76</sup> Although some studies have explored the impact of infliximab withdrawal on UC relapse, randomized trials assessing the withdrawal of mesalamine, thiopurines, non-TNF biologics, and tofacitinib have not been performed. In the absence of such data, decisions to discontinue treatment are typically made on a case-by-case basis. Recognizing that mesalamine can be discontinued in some patients who achieve remission on a biologic, Dr Abreu commented that there are others "who continue to have some symptoms even on a biologic who can benefit from oral and topical mesalamine." Additionally, it is reasonable to consider discontinuation of either thiopurines or biologics in patients who achieve deep remission on combination therapy, allowing

concentrations of each agent to guide the decision. For example, it may be appropriate to stop the thiopurine when infliximab levels are adequate, whereas infliximab may be discontinued in the setting of suboptimal infliximab concentrations but adequate thioguanine levels. Other patient factors, such as age and prognostic factors for relapse, may influence the decision as well. Regardless of which agent is discontinued, Dr Hanauer emphasized, "continued monitoring of symptoms, biomarkers, and endoscopy are necessary to avoid unanticipated disease progression."

## Updates on COVID-19 and IBD

### Overview of COVID-19

COVID-19, caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), was first identified in Wuhan, China, where over half of the 41 infected patients had direct contact history with a local seafood market selling live and slaughtered wild animals for food consumption.<sup>77-79</sup> Within months of the first reports in December 2019, the virus had infected thousands of people worldwide and had been declared a global pandemic.<sup>80</sup> At the time of this writing, nearly 18.7 million cases of COVID-19 and over 700,000 deaths have been confirmed worldwide.<sup>81</sup>

Named for its crown-like appearance,<sup>82</sup> SARS-CoV-2 is an enveloped RNA virus that enters target cells via the angiotensin-converting enzyme 2 (ACE2) receptor, which is expressed in epithelial cells of the lung, small intestine, colon, kidney, liver, pancreas, and blood vessels.<sup>83-86</sup> Transmission occurs primarily person-to-person via respiratory droplets from infected individuals and possibly from contaminated surfaces.<sup>87,88</sup> Manifestations of SARS-CoV-2 infection can range from asymptomatic to mild upper respiratory tract symptoms to life-threatening sepsis and cytokine storm. The most common manifestations among hospitalized patients with COVID-19

are fever and respiratory symptoms, including dry cough and shortness of breath.<sup>88,89</sup> Multiple reports have identified older age and the presence of comorbidities to be risk factors for severe COVID-19 and worse outcomes.<sup>77,90-94</sup>

### COVID-19 and the GI Tract

Although research conducted to date has focused on the mechanisms and consequences of SARS-CoV-2 infection in respiratory cells, the wide expression of ACE2 receptors throughout the digestive system, particularly the small intestine and colon, also provides a target for viral invasion.<sup>86,95,96</sup> Accordingly, it has become increasingly clear that a considerable

"Similar to the general population, patients with IBD should practice social distancing, appropriate use of personal protective equipment, and avoid nonessential travel to minimize potential exposure to other individuals with COVID-19."

– Dr Gary R. Lichtenstein

proportion of patients with COVID-19 experience gastrointestinal (GI) symptoms, usually in addition to respiratory symptoms.<sup>96-100</sup> An earlier report involving 204 patients with COVID-19 across hospitals in Hubei province in China found that half of patients reported a digestive symptom, most commonly lack of appetite and diarrhea.<sup>97</sup> Similarly, a multicenter cohort study across 9 US hospitals demonstrated that 61.3% of 318 patients with COVID-19 presented with at least 1 GI symptom, with lack

of appetite (34.8%), diarrhea (33.7%), and nausea (26.4%) being the most common symptoms.<sup>98</sup> In fact, GI symptoms were the chief complaint in 20% of patients and the initial presenting symptom in 14% of patients. Liver damage related to COVID-19 is also commonly observed, with a meta-analysis of 29 studies reporting a 15% to 20% pooled rate of elevated liver function enzymes among patients with COVID-19.<sup>96</sup>

A number of studies have detected SARS-CoV-2 RNA in stool specimens of infected patients, providing further evidence that the virus can involve the GI tract.<sup>86,89,95,96,100-103</sup> A recent meta-analysis found that 8 studies reported fecal tests that were positive for SARS-CoV-2, with fecal viral RNA shedding detected in 40.5% of patients.<sup>96</sup> Importantly, fecal viral shedding has been reported long after respiratory samples have become negative.<sup>101,103</sup> However, some investigators have found that the viral particles in stool were of insufficient quantities to be infectious.<sup>102</sup> Although these findings raise important questions, more data are needed to confirm the possibility of fecal-oral transmission and inform decisions regarding stool testing in infected patients.<sup>86,95,96,101</sup>

### **Risk, Clinical Course, and Outcomes of COVID-19 in IBD**

The potential for increased susceptibility of patients with IBD to SARS-CoV-2 infection and COVID-19 has been a concern due to the immune-mediated pathogenesis of the condition, frequent treatment with immunosuppressive therapies, and need to be at medical facilities for infusions or endoscopic procedures.<sup>104,105</sup> Although data are limited, one study reported a 0.0025 cumulative incidence of COVID-19 among 6000 patients with IBD in France and Italy, a rate similar to that observed in the general population.<sup>104</sup> Based on the limited published evidence and expert opinion, patients with IBD are not believed to have a baseline increased risk of infection with SARS-CoV-2 or development of

**Table 5.** Multivariate Regression for Outcomes From the SECURE-IBD Cohort

Variable (Referent)	ICU/Vent/Death OR (95% CI) N=517	P value
Age	1.04 (1.01-1.06)	.002
Male (female)	1.20 (0.55-2.60)	.65
Diagnosis of CD (UC/IBD unspecified)	0.76 (0.31-1.85)	.54
Active disease (remission)	1.14 (0.49-2.66)	.76
Systemic corticosteroid (none)	6.87 (2.30-20.51)	<.001
Anti-TNF agent (none)	0.90 (0.37-2.17)	.81
Current smoker	0.55 (0.06-4.94)	.59
BMI $\geq$ 30	2.00 (0.72-5.51)	.18
Comorbidities (none)		
1	1.22 (0.45-3.26)	.70
$\geq$ 2	2.87 (1.05-7.85)	.04
5-ASA/sulfasalazine (none)	3.14 (1.28-7.71)	.01

5-ASA, 5-aminosalicylate; BMI, body mass index; CD, Crohn's disease; IBD, inflammatory bowel disease; ICU, intensive care unit; OR, odds ratio; SECURE-IBD, Surveillance Epidemiology of Coronavirus Under Research Exclusion for Inflammatory Bowel Disease; TNF, tumor necrosis factor; UC, ulcerative colitis; vent, ventilator use. Adapted from Brenner et al.<sup>77</sup>

### COVID-19.<sup>105,106</sup>

Several reports have begun to characterize the clinical course and outcomes of COVID-19 in patients with IBD.<sup>77,90,107</sup> In a prospective observational cohort of patients with IBD and COVID-19 across 24 Italian referral units, 36 of 79 (46%) patients developed COVID-19–related pneumonia, 22 (28%) were hospitalized, 7 (9%) required nonmechanical ventilation, and 6 (8%) died.<sup>90</sup> Age over 65 years, active IBD, and higher Charleston Comorbidity Index score were significantly associated with worse prognosis (ie, higher risk of COVID-19–related pneumonia and death). Similar findings were observed in the first published analysis of data from the Surveillance Epidemiology of Coronavirus Under Research Exclusion for Inflammatory Bowel Disease (SECURE-IBD) registry.<sup>77</sup> In this report involving 525 cases from 33 countries, 37 (7%) patients developed severe COVID-19, 161 (31%) were hospitalized, and 16 died, translating to a 3% case fatality rate. Consistent with the Italian cohort, older age and presence of comorbidities were significantly associated with risk of

poor outcomes (composite outcome of ICU admission, ventilator support, and/or death) (Table 5). Although the presence of active disease significantly increased the risk of hospitalization or death, it was not associated with ICU admission/ventilator requirement/death. Taken collectively, these findings do not suggest that IBD predisposes patients to aggravated outcomes of COVID-19, and underscore the contribution of age and comorbidities to worse outcomes in IBD patients, as observed in the general population.

Given the increased risk of serious bacterial and viral infections associated with many IBD therapies,<sup>10,21,30,108,109</sup> the impact of these medications on the risk and clinical course of COVID-19 in patients with IBD has been an area of great interest.<sup>77,90,107</sup> To that end, a nationwide retrospective cohort study involving 36 of 37,857 IBD patients in the Veterans' Affairs Healthcare System with COVID-19 observed no increase of COVID-19 associated with thiopurine or anti-TNF therapy.<sup>107</sup> Further, published evidence to date has not demonstrated an association between worse outcomes of COVID-19 and

therapy with thiopurines, anti-TNF agents, or vedolizumab.<sup>77,90</sup> In contrast, evidence published to date has consistently implicated corticosteroids as a risk factor for worse outcomes in patients with IBD and COVID-19.<sup>77,90</sup> In the previously mentioned Italian study, use of corticosteroids was associated with increased risk of COVID-19–related pneumonia (OR, 4.94; 95% CI, 0.95-25.55;  $P=.05$ ) and death (OR, 6.28; 95% CI, 0.89-44.24;  $P=.064$ ).<sup>90</sup> Similarly, multivariate regression analyses of data from the SECURE-IBD registry found the use of systemic corticosteroids associated with an OR of 11.62 for death (95% CI, 2.09-64.74;  $P=.05$ ).<sup>77</sup> Of interest, 5-ASA/sulfasalazine use was also found to be significantly associated with worse outcomes in this population.


### Guidance for IBD Management During the Pandemic

In light of the potential implications of COVID-19 on patients with IBD, the American Gastroenterological Association has issued a clinical practice update summarizing available guidance for patients with IBD and their providers.<sup>105</sup> Recognizing that patients with IBD do not appear to be at higher risk of SARS-CoV-2 infection, patients who are not infected are encouraged to maintain their current regimens with the goal of maintaining clinical and endoscopic remission.<sup>105</sup> Despite concerns regarding patients going to infusion centers for delivery of certain therapies, the ongoing use of infusion centers is supported, provided that the center has appropriate COVID-19 screening protocols in place. Specific guidance for patients

who are infected with SARS-CoV-2 varies based on the presence and severity of COVID-19 manifestations, but general recommendations include tapering corticosteroids and holding thiopurines, methotrexate, and tofacitinib during the viral illness (Figure 7, Table 6). It is particularly important to investigate the source of GI symptoms in patients with COVID-19 and digestive symptoms, excluding enteric infection and active inflammation. Finally, clinicians who care for patients with IBD are encouraged to submit cases of IBD and confirmed COVID-19 to the SECURE-IBD registry at covidibd.org.

### Conclusion

The management of UC has evolved considerably over the past decade



**Figure 7.** American Gastroenterological Association guidance for managing patients with IBD and COVID-19. <sup>a</sup>Treatment of COVID-19 under investigation; consider therapies that have safety and efficacy in IBD. <sup>b</sup>Clearance of SARS-CoV-2 may enable resumption of IBD therapy; role of serologic antibody testing unclear at the current time. (Viral clearance testing may or may not be possible or appropriate, given local testing capabilities and health system–approved epidemiologic testing strategies during the COVID-19 pandemic.) 5-ASA, 5-aminosalicylate; COVID-19, coronavirus disease 2019; IBD, inflammatory bowel disease; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; SpO<sub>2</sub>, oxygen saturation. Adapted from Rubin et al.<sup>105</sup>



**Table 6.** American Gastroenterological Association Clinical Practice Update on Management of IBD During the COVID-19 Pandemic: Key Points

1. COVID-19 is the disease caused by the SARS-CoV-2 virus, but patients with IBD do not appear to be at a higher risk for infection with SARS-CoV-2 or development of COVID-19.
2. Patients with IBD who do not have infection with SARS-CoV-2 should not discontinue their IBD therapies and should continue infusion schedules at appropriate infusion centers.
3. Patients with IBD who have known SARS-CoV-2 but have not developed COVID-19 should hold thiopurines, methotrexate, and tofacitinib. Dosing of biologic therapies should be delayed for 2 weeks of monitoring for symptoms of COVID-19.
4. Patients with IBD who develop COVID-19 should hold thiopurines, methotrexate, tofacitinib, and biologic therapies during the viral illness. These can be restarted after complete symptom resolution or, if available, when follow-up viral testing is negative or serologic tests demonstrate the convalescent stage of illness.
5. The severity of the COVID-19 and the severity of the IBD should result in careful risk-benefit assessments regarding treatments for COVID-19 and escalating treatments for IBD.
6. Please submit cases of IBD and confirmed COVID-19 to the SECURE-IBD registry at covidibd.org.

COVID-19, coronavirus disease 2019; IBD, inflammatory bowel disease; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; SECURE-IBD, Surveillance Epidemiology of Coronavirus Under Research Exclusion for Inflammatory Bowel Disease. Adapted from Rubin et al.<sup>105</sup>

with growing emphasis on altering the natural history of the condition, recognition of both clinical and objective treatment targets, and availability of new treatment options. Taken collectively, these concepts have paved the way for a treat-to-target strategy in UC aimed at achieving and maintaining remission by adjusting therapies according to treatment targets. To that end, emerging data presented at DDW 2020 continue to add to our understanding of UC, strategies for optimizing its management, and the safety and efficacy of newer therapies. Lastly, clinicians who manage patients with UC are encouraged to keep up with the rapidly expanding evidence regarding COVID-19 and its implications for IBD management, and to report all cases of IBD and COVID-19 to the SECURE-IBD database.

## References

- Lo C-H, Lochhead P, Khalili H, et al. An inflammatory diet and risk of Crohn's disease and ulcerative colitis: a prospective cohort study [abstract 564]. *Gastroenterology*. 2020;158(6 suppl 1):S-116.
- Nguyen LH, Sikavi D, Ma W, et al. The sulfur microbial diet and risk of inflammatory bowel disease: results from three large prospective cohorts [abstract 119]. *Gastroenterology*. 2020;158(6 suppl 1):S-21.
- Bernstein CN, Walld R, Marrie RA. Social determinants of outcomes in IBD [abstract 26]. *Gastroenterology*. 2020;158(6 suppl 1):S-9-S-10.
- Nguyen GC, LaVeist TA, Harris ML, Wang MH, Datta LW, Brant SR. Racial disparities in utilization of specialist care and medications in inflammatory bowel disease. *Am J Gastroenterol*. 2010;105(10):2202-2208.
- Nguyen GC, Chong CA, Chong RY. National estimates of the burden of inflammatory bowel disease among racial and ethnic groups in the United States. *J Crohns Colitis*. 2014;8(4):288-295.
- Walker C, Allamneni C, Orr J, et al. Socioeconomic status and race are both independently associated with increased hospitalization rate among Crohn's disease patients. *Sci Rep*. 2018;8(1):4028.
- Kim ES, Tarassishin L, Eisele C, et al. Dynamics of intestinal inflammation and microbial dysbiosis in pregnant women with inflammatory bowel disease and their infants [abstract 423]. *Gastroenterology*. 2020;158(6 suppl 1):S-83-S-84.
- Khrom M, Long MD, Dube S, et al. Mega-analysis reveals novel genetic associations with extraintestinal manifestations in IBD [abstract 761]. *Gastroenterology*. 2020;158(6 suppl 1):S-157.
- Colombel JF, Panes J, D'Haens GR, et al. Higher versus standard adalimumab maintenance regimens in patients with moderately to severely active ulcerative colitis: results from the SERENE-UC maintenance study [abstract 945]. *Gastroenterology*. 2020;158(6 suppl 1):S-192.
- Humira [prescribing information]. North Chicago, IL: AbbVie; 2020.
- Yanai HA, Amir S, Avni-Biron I, et al. Increased risk of anti-drug antibodies to a second anti-TNF in patients who developed antibodies to the first anti-TNF [abstract Su1841]. *Gastroenterology*. 2020;158(6 suppl 1):S-671.
- Verstockt B, Outtier A, Lefrere J, et al. Endoscopic and histologic outcome in tofacitinib treated refractory moderate-to-severe ulcerative colitis: a prospective real-life cohort [abstract Mo1858]. *Gastroenterology*. 2020;158(6 suppl 1):S-952.
- Hu A, Kotze PG, Burgevin A, et al. Combination therapy does not improve rate of clinical or endoscopic remission in patients with inflammatory bowel diseases treated with vedolizumab or ustekinumab [published online July 12, 2020]. *Clin Gastroenterol Hepatol*. doi:10.1016/j.cgh.2020.07.012.
- Kobayashi T, Motoya S, Nakamura S, et al. The first prospective, multicenter, randomized controlled trial on discontinuation of maintenance infliximab in ulcerative colitis in remission: endoscopic normalization does not guarantee successful discontinuation [abstract 633]. *Gastroenterology*. 2020;158(6 suppl 1):S-136.
- Borren NZ, Plichta D, Joshi AD, et al. Multi-omics profiling in patients with quiescent inflammatory bowel disease identifies biomarkers predicting relapse [abstract 348]. *Gastroenterology*. 2020;158(6 suppl 1):S-63.
- Waterman M, Mendelovici A, Harel S, et al. Association of lymphoma with age at onset and thiopurine use among patients with inflammatory bowel diseases—results from the epidemiology group of the Israeli IBD Research Nucleus [abstract 850]. *Gastroenterology*. 2020;158(6 suppl 1):S-172-S-173.
- Khan N, Abbas AM, Lichtenstein GR, Loftus EV Jr, Bazzano LA. Risk of lymphoma in patients with ulcerative colitis treated with thiopurines: a nationwide retrospective cohort study. *Gastroenterology*. 2013;145(5):1007-1015.e3.
- Kotlyar DS, Lewis JD, Beaugerie L, et al. Risk of lymphoma in patients with inflammatory bowel disease treated with azathioprine and 6-mercaptopurine: a meta-analysis. *Clin Gastroenterol Hepatol*. 2015;13(5):847-858.e4.
- Lemaitre M, Kirchgessner J, Rudnichi A, et al. Association between use of thiopurines or tumor necrosis factor antagonists alone or in combination and risk of lymphoma in patients with inflammatory bowel disease. *JAMA*. 2017;318(17):1679-1686.
- Patel D, Kavani H, Trivedi C, Pernes T, Khan NH. The incidence of pneumonia and impact of immunosuppressive medications on risk of pneumonia among patients with IBD [abstract 282]. *Gastroenterology*. 2020;158(6 suppl 1):S-54-S-55.
- Xeljanz [prescribing information]. NY, NY: Pfizer Labs; 2019.
- Sandborn WJ, Panes J, Sands BE, et al. Incidence of venous thromboembolic events in patients with ulcerative colitis treated with tofacitinib in the ulcerative colitis clinical development program: an update as of May 2019 [abstract 684]. *Gastroenterology*. 2020;158(6 suppl 1):S-144.
- Sandborn WJ, Panes J, Sands BE, et al. Venous thromboembolic events in the tofacitinib ulcerative colitis clinical development programme. *Aliment Pharmacol Ther*. 2019;50(10):1068-1076.
- Goettl JA, Kotlarz D, Emani R, et al. Low-dose interleukin-2 ameliorates colitis in a preclinical humanized mouse model. *Cell Mol Gastroenterol Hepatol*. 2019;8(2):193-195.
- Allegretti JR, Canavan J, Mitsialis V, et al. Low dose interleukin-2 for the treatment of moderate to severe ulcerative colitis [abstract 1026]. *Gastroenterology*. 2020;158(6 suppl 1):S-206.
- Reinisch W, Leszczyszyn J, Dudkowiak R, et al. A novel subcutaneous infliximab (CT-P13) in patients with active Crohn's disease and ulcerative colitis: week 54 and switching results from a multicenter, randomised controlled pivotal trial [abstract 948]. *Gastroenterology*. 2020;158(6 suppl 1):S-193.
- Colombel JF, D'Haens G, Lee WJ, Petersson J, Panaccione R. Outcomes and strategies to support a

- treat-to-target approach in inflammatory bowel disease: a systematic review. *J Crohns Colitis*. 2020;14(2):254-266.
28. Peyrin-Biroulet L, Sandborn W, Sands BE, et al. Selecting Therapeutic Targets in Inflammatory Bowel Disease (STRIDE): determining therapeutic goals for treat-to-target. *Am J Gastroenterol*. 2015;110(9):1324-1338.
29. Ungaro R, Colombel J-F, Lissos T, Peyrin-Biroulet L. A treat-to-target update in ulcerative colitis: a systematic review. *Am J Gastroenterol*. 2019;114(6):874-883.
30. Rubin DT, Ananthakrishnan AN, Siegel CA, Sauer BG, Long MD. ACG Clinical Guideline: ulcerative colitis in adults. *Am J Gastroenterol*. 2019;114(3):384-413.
31. Rosenberg L, Lawlor GO, Zenlea T, et al. Predictors of endoscopic inflammation in patients with ulcerative colitis in clinical remission. *Inflamm Bowel Dis*. 2013;19(4):779-784.
32. Eriksson C, Cao Y, Rundquist S, et al. Changes in medical management and colectomy rates: a population-based cohort study on the epidemiology and natural history of ulcerative colitis in Örebro, Sweden, 1963-2010. *Aliment Pharmacol Ther*. 2017;46(8):748-757.
33. Boal Carvalho P, Cotter J. Mucosal healing in ulcerative colitis: a comprehensive review. *Drugs*. 2017;77(2):159-173.
34. Battar R, Duijvestein M, Guizzetti L, et al. Histologic healing rates of medical therapies for ulcerative colitis: a systematic review and meta-analysis of randomized controlled trials. *Am J Gastroenterol*. 2019;114(5):733-745.
35. Bryant RV, Winer S, Travis SP, Riddell RH. Systematic review: histological remission in inflammatory bowel disease. Is 'complete' remission the new treatment paradigm? An IOIBD initiative. *J Crohns Colitis*. 2014;8(12):1582-1597.
36. Marchal Bressenot A, Riddell RH, Boulaignon-Rombi C, et al. Review article: the histological assessment of disease activity in ulcerative colitis. *Aliment Pharmacol Ther*. 2015;42(8):957-967.
37. Peyrin-Biroulet L, Bressenot A, Kampman W. Histologic remission: the ultimate therapeutic goal in ulcerative colitis? *Clin Gastroenterol Hepatol*. 2014;12(6):929-934.e2.
38. Pai RK, Jairath V, Vande Casteele N, Rieder F, Parker CE, Lauwers GY. The emerging role of histologic disease activity assessment in ulcerative colitis. *Gastrointest Endosc*. 2018;88(6):887-898.
39. Colman RJ, Rubin DT. Histological inflammation increases the risk of colorectal neoplasia in ulcerative colitis: a systematic review. *Intest Res*. 2016;14(3):202-210.
40. Rubin DT, Huo D, Kinnucan JA, et al. Inflammation is an independent risk factor for colonic neoplasia in patients with ulcerative colitis: a case-control study. *Clin Gastroenterol Hepatol*. 2013;11(12):1601-1608.e1-e4.
41. Alper A, Zhang L, Pashankar DS. Correlation of erythrocyte sedimentation rate and C-reactive protein with pediatric inflammatory bowel disease activity. *J Pediatr Gastroenterol Nutr*. 2017;65(2):e25-e27.
42. Fagan EA, Dyck RF, Maton PN, et al. Serum levels of C-reactive protein in Crohn's disease and ulcerative colitis. *Eur J Clin Invest*. 1982;12(4):351-359.
43. Ananthakrishnan AN. Epidemiology and risk factors for IBD. *Nat Rev Gastroenterol Hepatol*. 2015;12(4):205-217.
44. Regueiro MD, Greer JB, Hanauer SB. Established management paradigms in IBD: treatment targets and therapeutic tools. *Am J Gastroenterol Suppl*. 2016;3:8-16.
45. D'Haens GR, Sartor RB, Silverberg MS, Petersson J, Rutgeerts P. Future directions in inflammatory bowel disease management. *J Crohns Colitis*. 2014;8(8):726-734.
46. Dassopoulos T, Cohen RD, Scherl EJ, Schwartz RM, Kosinski L, Regueiro MD. Ulcerative colitis care pathway. *Gastroenterology*. 2015;149(1):238-245.
47. Feuerstein JD, Isaacs KL, Schneider Y, Siddique SM, Falck-Ytter Y, Singh S; AGA Institute Clinical Guidelines Committee. AGA Clinical Practice Guidelines on the management of moderate to severe ulcerative colitis. *Gastroenterology*. 2020;158(5):1450-1461.
48. Singh S, Fumery M, Sandborn WJ, Murad MH. Systematic review with network meta-analysis: first- and second-line pharmacotherapy for moderate-severe ulcerative colitis. *Aliment Pharmacol Ther*. 2018;47(2):162-175.
49. Feagan BG, Reinisch W, Rutgeerts P, et al. The effects of infliximab therapy on health-related quality of life in ulcerative colitis patients. *Am J Gastroenterol*. 2007;102(4):794-802.
50. Ford AC, Sandborn WJ, Khan KJ, Hanauer SB, Talley NJ, Moayyedi P. Efficacy of biological therapies in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol*. 2011;106(4):644-659.
51. Rutgeerts P, Sandborn WJ, Feagan BG, et al. Infliximab for induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2005;353(23):2462-2476.
52. Sandborn WJ, Rutgeerts P, Feagan BG, et al. Colectomy rate comparison after treatment of ulcerative colitis with placebo or infliximab. *Gastroenterology*. 2009;137(4):1250-1260.
53. Sandborn WJ, Feagan BG, Marano C, et al; PURSUIT-Maintenance Study Group. Subcutaneous golimumab maintains clinical response in patients with moderate-to-severe ulcerative colitis. *Gastroenterology*. 2014;146(1):96-109.e1.
54. Sandborn WJ, Feagan BG, Marano C, et al; PURSUIT-SC Study Group. Subcutaneous golimumab induces clinical response and remission in patients with moderate-to-severe ulcerative colitis. *Gastroenterology*. 2014;146(1):85-95.
55. Hindryckx P, Vande Casteele N, Novak G, et al. The expanding therapeutic armamentarium for inflammatory bowel disease: how to choose the right drug(s) for our patients. *J Crohns Colitis*. 2018;12(1):105-119.
56. Arora Z, Shen B. Biological therapy for ulcerative colitis. *Gastroenterol Rep (Oxf)*. 2015;3(2):103-109.
57. Pouillon L, Van Stappen J, Bossuyt P, Danese S, Peyrin-Biroulet L. Should we use anti-tumor necrosis factor agents or vedolizumab as first-line biological therapy in ulcerative colitis? *Best Pract Res Clin Gastroenterol*. 2018;32:33-17-25.
58. Colombel J-F, Sands BE, Rutgeerts P, et al. The safety of vedolizumab for ulcerative colitis and Crohn's disease. *Gut*. 2017;66(5):839-851.
59. Sandborn WJ, Su C, Sands BE, et al; OCTAVE Induction 1, OCTAVE Induction 2, and OCTAVE Sustain Investigators. Tofacitinib as induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2017;376(18):1723-1736.
60. Berinstein JA, Steiner CA, Regal RE, et al. Efficacy of induction therapy with high-intensity tofacitinib in 4 patients with acute severe ulcerative colitis. *Clin Gastroenterol Hepatol*. 2019;17(5):988-990.e1.
61. Panaccione R, Ghosh S, Middleton S, et al. Combination therapy with infliximab and azathioprine is superior to monotherapy with either agent in ulcerative colitis. *Gastroenterology*. 2014;146(2):392-400.e3.
62. Afif W, Loftus EV Jr, Faubion WA, et al. Clinical utility of measuring infliximab and human anti-chimeric antibody concentrations in patients with inflammatory bowel disease. *Am J Gastroenterol*. 2010;105(5):1133-1139.
63. Baert F, Noman M, Vermeire S, et al. Influence of immunogenicity on the long-term efficacy of infliximab in Crohn's disease. *N Engl J Med*. 2003;348(7):601-608.
64. Maser EA, Vilella R, Silverberg MS, Greenberg GR. Association of trough serum infliximab to clinical outcome after scheduled maintenance treatment for Crohn's disease. *Clin Gastroenterol Hepatol*. 2006;4(10):1248-1254.
65. Papamichael K, Cheifetz AS. Use of anti-TNF drug levels to optimise patient management. *Frontline Gastroenterol*. 2016;7(4):289-300.
66. Billioud V, Sandborn WJ, Peyrin-Biroulet L. Loss of response and need for adalimumab dose intensification in Crohn's disease: a systematic review. *Am J Gastroenterol*. 2011;106(4):674-684.
67. Hanauer SB, Wagner CL, Bala M, et al. Incidence and importance of antibody responses to infliximab after maintenance or episodic treatment in Crohn's disease. *Clin Gastroenterol Hepatol*. 2004;2(7):542-553.
68. Feagan BG, Rutgeerts P, Sands BE, et al; GEMINI 1 Study Group. Vedolizumab as induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2013;369(8):699-710.
69. Pouillon L, Vermeire S, Bossuyt P. Vedolizumab trough level monitoring in inflammatory bowel disease: a state-of-the-art overview. *BMC Med*. 2019;17(1):89.
70. Adedokun OJ, Xu Z, Marano C, et al. Ustekinumab pharmacokinetics and exposure responses in a phase 3 randomized trial of patients with ulcerative colitis. *Clin Gastroenterol Hepatol*. 2020;18(10):2244-2255.e9.
71. Feuerstein JD, Nguyen GC, Kupfer SS, Falck-Ytter Y, Singh S; American Gastroenterological Association Institute Clinical Guidelines Committee. American Gastroenterological Association Institute Guideline on therapeutic drug monitoring in inflammatory bowel disease. *Gastroenterology*. 2017;153(3):827-834.
72. Papamichael K, Baert F, Tops S, et al. Post-induction adalimumab concentration is associated with short-term mucosal healing in patients with ulcerative colitis. *J Crohns Colitis*. 2017;11(1):53-59.
73. Vande Casteele N, Ferrante M, Van Assche G, et al. Trough concentrations of infliximab guide dosing for patients with inflammatory bowel disease. *Gastroenterology*. 2015;148(7):1320-1329.e3.
74. de Boer NKH, Peyrin-Biroulet L, Jharap B, et al. Thiopurines in inflammatory bowel disease: new findings and perspectives. *J Crohns Colitis*. 2018;12(5):610-620.
75. Fiorino G, Cortes PN, Ellul P, et al. Discontinuation of infliximab in patients with ulcerative colitis is associated with increased risk of relapse: a multinational retrospective cohort study. *Clin Gastroenterol Hepatol*. 2016;14(10):1426-1432.e1.
76. Kennedy NA, Warner B, Johnston EL, et al; UK Anti-TNF Withdrawal Study Group. Relapse after withdrawal from anti-TNF therapy for inflammatory bowel disease: an observational study, plus systematic review and meta-analysis. *Aliment Pharmacol Ther*. 2016;43(8):910-923.
77. Brenner EJ, Ungaro RC, Geary RB, et al. Corticosteroids, but not TNF antagonists, are associated with adverse COVID-19 outcomes in patients with inflammatory bowel disease: results from an international registry. *Gastroenterology*. 2020;159(2):481-491.e3.
78. Xu Y. Unveiling the origin and transmission of 2019-nCoV. *Trends Microbiol*. 2020;28(4):239-240.
79. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-273.
80. World Health Organization. WHO Direc-

- tor-General's opening remarks at the media briefing on COVID-19—11 March 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. Accessed July 26, 2020.
81. Johns Hopkins Coronavirus Resource Center. <https://coronavirus.jhu.edu>. Accessed July 26, 2020.
82. Ma RCW, Holt RIG. COVID-19 and diabetes. *Diabet Med*. 2020;37(5):723-725.
83. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol*. 2020;5(4):562-569.
84. Walls AC, Park Y-J, Tortorici MA, Wall A, McGuire AT, Veers D. Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell*. 2020;181(2):281-292.e6.
85. Muniyappa R, Gubbi S. COVID-19 pandemic, coronaviruses, and diabetes mellitus. *Am J Physiol Endocrinol Metab*. 2020;318(5):E736-E741.
86. Ma C, Cong Y, Zhang H. COVID-19 and the digestive system. *Am J Gastroenterol*. 2020;115(7):1003-1006.
87. van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med*. 2020;382(16):1564-1567.
88. Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): a review [published online July 10, 2020]. *JAMA*. doi:10.1001/jama.2020.12839.
89. Mao R, Qiu Y, He JS, et al. Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol*. 2020;5(7):667-678.
90. Bezzio C, Saibeni S, Variola A, et al; Italian Group for the Study of Inflammatory Bowel Disease (IG-IBD). Outcomes of COVID-19 in 79 patients with IBD in Italy: an IG-IBD study. *Gut*. 2020;69(7):1213-1217.
91. Gold MS, Sehayek D, Gabrielli S, Zhang X, McCusker C, Ben-Shoshan M. COVID-19 and comorbidities: a systematic review and meta-analysis [published online July 14, 2020]. *Postgrad Med*. doi:10.1080/00325481.2020.1786964.
92. Guan WJ, Ni ZY, Hu Y, et al; China Medical Treatment Expert Group for Covid-19. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-1720.
93. Guan WJ, Liang WH, Zhao Y, et al; China Medical Treatment Expert Group for COVID-19. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J*. 2020;55(5):2000547.
94. Wu D, Wu T, Liu Q, Yang Z. The SARS-CoV-2 outbreak: what we know. *Int J Infect Dis*. 2020;94:44-48.
95. Ng SC, Tilg H. COVID-19 and the gastrointestinal tract: more than meets the eye. *Gut*. 2020;69(6):973-974.
96. Parasa S, Desai M, Thoguluva Chandrasekar V, et al. Prevalence of gastrointestinal symptoms and fecal viral shedding in patients with coronavirus disease 2019. A systematic review and meta-analysis. *JAMA Netw Open*. 2020;3(6):e2011335.
97. Pan L, Mu M, Yang P, et al. Clinical characteristics of COVID-19 patients with digestive symptoms in Hubei, China: a descriptive, cross-sectional, multicenter study. *Am J Gastroenterol*. 2020;115(5):766-773.
98. Redd WD, Zhou JC, Hathorn KE, et al. Prevalence and characteristics of gastrointestinal symptoms in patients with severe acute respiratory syndrome coronavirus 2 infection in the United States: a multicenter cohort study. *Gastroenterology*. 2020;159(2):765-767.e2.
99. Tariq R, Saha S, Furqan F, Hassett L, Pardi D, Khanna S. Prevalence and mortality of COVID-19 patients with gastrointestinal symptoms: a systematic review and meta-analysis. *Mayo Clin Proc*. 2020;95(8):1632-1648.
100. Zhou Z, Zhao N, Shu Y, Han S, Chen B, Shu X. Effect of gastrointestinal symptoms on patients infected with COVID-19. *Gastroenterology*. 2020;158(8):2294-2297.
101. Wu Y, Guo C, Tang L, et al. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol*. 2020;5(5):434-435.
102. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature*. 2020;581(7809):465-469.
103. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology*. 2020;158(6):1831-1833.e3.
104. Allocca M, Fiorino G, Zallot C, et al. Incidence and patterns of COVID-19 among inflammatory bowel disease patients from the Nancy and Milan cohorts. *Clin Gastroenterol Hepatol*. 2020;18(9):2134-2135.
105. Rubin DT, Feuerstein JD, Wang AY, Cohen RD. AGA Clinical Practice Update on management of inflammatory bowel disease during the COVID-19 pandemic: expert commentary. *Gastroenterology*. 2020;159(1):350-357.
106. Neurath MF. COVID-19 and immunomodulation in IBD. *Gut*. 2020;69(7):1335-1342.
107. Khan N, Patel D, Xie D, Lewis J, Trivedi C, Yang YX. Impact of anti-TNF and thiopurine medications on the development of COVID-19 in patients with inflammatory bowel disease: a nationwide VA cohort study [published online May 29, 2020]. *Gastroenterology*. doi:10.1053/j.gastro.2020.05.065.
108. Lichtenstein GR, Loftus EV, Isaacs KL, Regueiro MD, Gerson LB, Sands BE. ACG Clinical Guideline: management of Crohn's disease in adults. *Am J Gastroenterol*. 2018;113(4):481-517.
109. Ford AC, Peyrin-Biroulet L. Opportunistic infections with anti-tumor necrosis factor- $\alpha$  therapy in inflammatory bowel disease: meta-analysis of randomized controlled trials. *Am J Gastroenterol*. 2013;108(8):1268-1276.

